

# Tobacco Budworm



**T**he tobacco budworm is an agricultural pest moth species that attacks agricultural crops and ornamental flowers. The larvae, or caterpillars, feed on the buds, flowers, and fruits of host plants, causing damage.

Tobacco Budworm moths emerge from overwintering pupae from March through May and lay eggs of Petunia, Geranium, Nicotiana, and snapdragon. Corn Earworms, which have a similar appearance, will target corn and lettuce.

The larvae are predominantly yellow green but have the ability to take on the hue of whatever flower they happen to be eating. This camouflage is so good that they are usually not noticed until the affected plant is no longer blooming as the larvae feed on foliage as well as seed capsules.

Caterpillars can be controlled by handpicking or by natural predators such as the egg parasitizing Trichogramma wasp. Other insects that eat the eggs and small larvae are the minute pirate bug and bigeyed bug.

If spraying is necessary, an organic control such as Monterey Bt Caterpillar Killer is recommended. Spinosad products such as Bonide Captain Jack's could also be used but will harm beneficial insects and can only be applied 3x a season to the same crop or plant. Spinosad is especially harmful to syrphid fly larvae that help control corn earworm eggs and small larvae on lettuce.



**We'll help you grow the plants you love!**

[www.sloatgardens.com](http://www.sloatgardens.com)